

Recombinant Bcl-6 (Follicular Lymphoma Marker) Antibody

Rabbit Monoclonal Antibody [Clone BCL6/8983R]

Catalog No	Format	Size
604-RBM30-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
604-RBM30-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
604-RBM30-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	BCL6/8983R
Gene Name	BCL6
Immunogen	Recombinant fragment (aa1-400) of human BCL6 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	95kDa
Cellular Localization	Nucleus.
Species Reactivity	Human
Positive Control	Raji or Ramos cells. Human tonsil or Hodgkin s lymphoma.


*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Bcl-6 (Follicular Lymphoma Marker) Antibody



SDS-PAGE Analysis of Purified bcl-6 Recombinant Rabbit Monoclonal Antibody (BCL6/8983R). Confirmation of Purity and Integrity of Antibody.

Western Blot Analysis of Raji and Ramos lysates using BCL6 Recombinant Rabbit Monoclonal Antibody (BCL6/8983R).



Formalin-fixed, paraffin-embedded human tonsil stained with BCL-6 Rabbit Recombinant Monoclonal Antibody (BCL6/8983R). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large (L&H) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Cytokine Signaling, Immunology, Nuclear Marker, Transcription Factors