

ERCC1 / RAD10 (Tumor Progression Marker) Antibody

Mouse Monoclonal Antibody [Clone ERCC1/7597]

Catalog No	Format	Size
2067-MSM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2067-MSM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2067-MSM6-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	ERCC1/7597
Gene Name	ERCC1
Immunogen	Recombinant fragment (around aa 191-281) of human ERCC1 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	110kDa
Cellular Localization	Nucleus.
Species Reactivity	Human
Positive Control	HeLa or HepG2 cells. Human tonsil or prostate. Brain, Pancreas or Spleen.

*Optimal dilution for a specific application should be determined.


Product Images for ERCC1 / RAD10 (Tumor Progression Marker) Antibody



SDS-PAGE Analysis of Purified ERCC1 Mouse Monoclonal Antibody (ERCC1/7597). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Brain, Human Pancreas and Human Spleen tissue lysates using ERCC1 Mouse Monoclonal Antibody (ERCC1/7597).



Formalin-fixed, paraffin-embedded human bladder stained with ERCC1 Mouse Monoclonal Antibody (ERCC1/7597). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Recognizes a protein of 110kDa, identified as Excision Repair Cross Complementing 1 (ERCC1). It is a mammalian nucleotide excision repair (NER) enzyme involved in repair of damaged DNA. ERCC1 is a homologous to RAD10 in *Saccharomyces cerevisiae*, which is required in mitotic intrachromosomal recombination and repair. ERCC1 is required in repair of cisplatin-induced DNA adducts and ultraviolet (UV)-induced DNA damage. High expression of ERCC1 has been linked to tumor progression in a variety of cancers including non-small cell lung cancer (NSCLC), squamous cell carcinoma of the head, ovarian cancer and esophageal cancer.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Nuclear Marker
