

IL1RN / IL-1RN / IL1RA (Interleukin-1RA) Antibody

Mouse Monoclonal Antibody [Clone IL1RA/4715]


Catalog No	Format	Size
3557-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3557-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3557-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes


Product Details	
Clone	IL1RA/4715
Gene Name	IL1RN
Immunogen	Recombinant fragment (around aa1-200) of human Interleukin-1RA protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	17-25kDa
Cellular Localization	Cytoplasm. Secreted.
Species Reactivity	Human
Positive Control	Epithelial cells.

*Optimal dilution for a specific application should be determined.


Product Images for IL1RN / IL-1RN / IL1RA (Interleukin-1RA) Antibody



Formalin-fixed, paraffin-embedded human tonsil stained with Interleukin-1RA Mouse Monoclonal Antibody (IL1RA/4715). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



Formalin-fixed, paraffin-embedded human adrenal gland stained with Interleukin-1RA Mouse Monoclonal Antibody (IL1RA/4715). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.




Analysis of Protein Array containing more than 19,000 full-length human proteins using Interleukin-1RA Mouse Monoclonal Antibody (IL1RA/4715). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

Two forms of interleukin-1, designated IL-1 α and IL-1 β , have been described. Although encoded by distinct genes and exhibiting roughly only 25% sequence identity, IL-1 α and IL-1 β bind to the same receptor and seem to elicit similar biological responses. IL-1 production is generally thought to be associated with inflammation, but it has also been shown to be expressed during kidney development, thymocyte differentiation and cartilage degradation. IL-1 plays a critical role in the regulation of immune response and inflammation acting as an activator of T and B lymphocytes and natural killer (NK) cells. IL-1 receptor antagonist (IL-1ra) is a cytokine that inhibits IL-1 α and IL-1 β binding to interleukin receptors. By neutralizing the activity of IL-1, IL-1ra contributes to the inhibition of the immune and inflammatory responses and has been targeted as a drug for the treatment of severely active rheumatoid arthritis. There are four isoforms of IL-1ra that are produced as a result of alternative splicing events.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.



SDS-PAGE Analysis of Purified Interleukin-1RA Mouse Monoclonal Antibody (IL1RA/4715). Confirmation of Purity and Integrity of Antibody.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Cytokine Signaling, Immunology, Nuclear Marker