

Recombinant CD163 (Monocyte & Macrophage Marker) Antibody

Mouse Monoclonal Antibody [Clone r1032]

Catalog No	Format	Size
9961-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
9961-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
9961-MSM4-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	r1032
Immunogen	Recombinant full-length human Major Vault Protein (MVP) protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	99.33kDa
Cellular Localization	Cytoplasm, Nuclear pore complex, Nucleus, Perinuclear region
Species Reactivity	Human
Positive Control	MCF-7 or HeLa cells. Human breast tumors. Lung, Spleen, Stomach and A549

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant CD163 (Monocyte & Macrophage Marker) Antibody



Formalin-fixed, paraffin-embedded human prostate stained with MVP Recombinant Mouse Monoclonal Antibody (r1032).



SDS-PAGE Analysis of Purified MVP Recombinant Mouse Monoclonal Antibody (r1032). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Lung, Human Spleen tissue, Human Stomach and A549 lysates using Major Vault Protein Recombinant Rabbit Monoclonal Antibody (r1032).

Specificity & Comments

Recognizes a protein of 104kDa-110kDa, characterized as major vault protein (MVP). Vaults are large ribonucleoprotein particles (RNPs) present in all eukaryotic cells. They have a complex morphology, including several small molecules of RNA, but a single protein species. The MVP accounts for >70% of their mass. Their shape is reminiscent of the nucleopore central plug. Treatment of cells with estradiol increases the amount of MVP in nuclear extract. The hormone-dependent interaction of vaults with ER is prevented in vitro by sodium molybdate. Antibodies to estrogen, progesterone and glucocorticoid receptors are able to co-immunoprecipitate the MVP. MVP is overexpressed in many neoplastic tissues and cell lines. Expression of MVP predicts a poor response to chemotherapy.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.