

## Recombinant CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone NCAM1/9074R]

Catalog No	Format	Size
4684-RBM36-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4684-RBM36-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4684-RBM36-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	NCAM1/9074R
<b>Gene Name</b>	NCAM1
<b>Immunogen</b>	Recombinant fragment of human NCAM1 (CD56) protein (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	145 and 125kDa 180
<b>Cellular Localization</b>	Cell Surface. Cytoplasm.
<b>Species Reactivity</b>	Human, Mouse
<b>Positive Control</b>	Human colon or small intestine. Human Brain, Mouse Brain

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Antibody




SDS-PAGE Analysis of Purified CD56 Rabbit Recombinant Monoclonal Antibody (NCAM1/9074R). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of Human Brain tissue lysates using CD56 Rabbit Recombinant Monoclonal Antibody (NCAM1/9074R).



Western blot analysis of Human Brain, Mouse Brain and Rat Brain tissue lysates using CD56 Rabbit Recombinant Monoclonal Antibody (NCAM1/9074R).



Formalin-fixed, paraffin-embedded human colon stained with CD56 Rabbit Recombinant Monoclonal Antibody (NCAM1/9074R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

### Specificity & Comments

This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Cytokine Signaling, Developmental Biology, Hematopoietic Stem Cells, Immunology, Mesenchymal Stem Cell Differentiation, Neural Stem Cells, Neuroscience, Signal Transduction, Stem Cell Differentiation

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.