

RET Proto-oncogene Antibody

Mouse Monoclonal Antibody [Clone RET/7694]

Catalog No	Format	Size
5979-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5979-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5979-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	RET/7694
Gene Name	RET
Immunogen	Recombinant fragment (around aa 702-848) of human RET protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	150kDa (precursor); 170kDa (Mature)
Cellular Localization	Cytoplasm. Cell Surface. Nucleus.
Species Reactivity	Human
Positive Control	Human breast prostate or colon carcinoma. Brain.

*Optimal dilution for a specific application should be determined.


Product Images for RET Proto-oncogene Antibody



SDS-PAGE Analysis of Purified RET Proto-oncogene Mouse Monoclonal Antibody (RET/7694). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Brain tissue lysate using RET Oncogene Mouse Monoclonal Antibody (RET/7694).



Formalin-fixed, paraffin-embedded human parathyroid gland stained with RET Proto-oncogene Mouse Monoclonal Antibody (RET/7694). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

The Ret proto-oncogene is structurally related to the growing family of tyrosine kinase transmembrane receptors and is involved in GDNF signaling. RET expression is reported in several regions of the central nervous system; in the developing cranial nerve ganglia and a subset of cells within dorsal root ganglia, in motor neurons in the spinal cord and hindbrain, in neuro-retina and the growing tips of the renal collecting ducts in developing kidney. Alterations in RET gene are associated with diseases including papillary thyroid carcinoma, multiple endocrine neoplasia (type 2A and 2B), familial medullary thyroid carcinoma, and a congenital developmental disorder known as Hirschsprung's disease.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Neuroscience, Signal Transduction, Transcription Factors