

Recombinant Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone rKRTH/8883]

Catalog No	Format	Size
MSM58-8883-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM58-8883-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM58-8883-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rKRTH/8883
Gene Name	KRT76
Immunogen	Recombinant full-length human KRT76 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	52-67kDa
Cellular Localization	Cytoplasm.
Species Reactivity	Human
Positive Control	Epithelial cells Skin or Adenocarcinomas, Skin, A431.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody



SDS-PAGE Analysis of Purified CK HMW Recombinant Mouse Monoclonal Antibody (rKRTH/8883). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Skin and A431 lysates using CytokeratinHMW Recombinant Mouse Monoclonal Antibody (rKRTH/8883).



Formalin-fixed, paraffin-embedded human skin stained with Cytokeratin, HMW Recombinant Mouse Monoclonal Antibody (rKRTH/8883). HIER: Tris/EDTA, pH9.0, 45min. 2°C; HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. The acidic keratins have molecular weights (MW) of 56.5, 55, 51, 50, 50, 48, 46, 45, and 40kDa. MAb AE3 recognizes the 65-67, 64, 59, 58, 56, and 52kDa keratins of basic subfamily. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology
