

Recombinant Clusterin / Apolipoprotein J (APO-J) Antibody

Rabbit Monoclonal Antibody [Clone CLU/8865R]

Catalog No	Format	Size
1191-RBM24-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1191-RBM24-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1191-RBM24-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	CLU/8865R
Gene Name	CLU
Immunogen	Recombinant fragment (around aa410-440) of human CLU protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	70kDa (precursor); 36-39kDa ($\hat{\mu}$); 34-36kDa ($\hat{\mu}$ 2)
Cellular Localization	Cytoplasm, Secreted
Species Reactivity	Human
Positive Control	Human pancreas tonsil or cervix.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Clusterin / Apolipoprotein J (APO-J) Antibody



SDS-PAGE Analysis of Purified Clusterin | Clusterin beta chain | Clusterin alpha chain Recombinant Rabbit Monoclonal Antibody (CLU/8865R). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human tonsil stained with Clusterin / APOJ Recombinant Rabbit Monoclonal Antibody (CLU/8865R). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Clusterin, also designated complement lysis inhibitor (CLI), apolipoprotein J (APOJ), sulfated glycoprotein 2 (SGP2), SP40 and testosterone-repressed prostate message 2 (TRPM2), is a secretory, heterodimeric glycoprotein that influences immune regulation, cell adhesion, transformation, lipid transportation, tissue remodeling, membrane recycling and cell-cell interactions. Clusterin is synthesized as a 449 amino acid polypeptide that is post-translationally cleaved at an internal bond between Arg 227 and Ser 228. Two subunits, ? and ?, are associated through disulfide bonds. The ? subunit (also called ApoJ?) corresponds to residues 23-227. The ? subunit (also called ApoJ?) corresponds to residues 228-449. Overexpression of Clusterin appears to be more common in late stages of mammary tumor progression. Clusterin markedly influences ?-Amyloid structure and neuritic toxicity in vivo and may influence Alzheimer's pathogenesis.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Neuroscience, Complement System