

Recombinant Fatty Acid Binding Protein (Liver) / FABP1 Antibody

Rabbit Monoclonal Antibody [Clone FABP1/9085R]

Catalog No	Format	Size
2168-RBM26-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2168-RBM26-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2168-RBM26-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	FABP1/9085R
Immunogen	Recombinant fragment (around aa1-127) of the human FABP1 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	14.21kDa
Cellular Localization	Cytoplasm
Species Reactivity	Human
Positive Control	Liver or colon carcinoma tissues (IHC). Human kidney tissue lysate (WB). intestine.



*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Fatty Acid Binding Protein (Liver) / FABP1 Antibody




Formalin-fixed, paraffin-embedded human kidney stained with FABP1 Recombinant Rabbit Monoclonal Antibody (FABP1/9085R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

SDS-PAGE Analysis of Purified FABP1 Recombinant Rabbit Monoclonal Antibody (FABP1/9085R). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of human intestine using FABP1 Recombinant Rabbit Monoclonal Antibody (FABP1/9085R).

Western blot analysis of intestine tissue lysates of different species using FABP1 Recombinant Rabbit Monoclonal Antibody (FABP1/9085R).



Formalin-fixed, paraffin-embedded human kidney stained with FABP1 Recombinant Rabbit Monoclonal Antibody (FABP1/9085R). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Fatty acid-binding proteins, designated FABPs, are a family of homologous cytoplasmic proteins that are expressed in a highly tissue-specific manner and play an integral role in the balance between lipid and carbohydrate metabolism. FABPs mediate fatty acid (FA) and/or hydrophobic ligand uptake, transport and targeting within their respective tissues. The mechanisms underlying these actions can give rise to both passive diffusional uptake and protein-mediated transmembrane transport of FAs. FABPs are expressed in adipocytes (A-FABP), brain (B-FABP), epithelium (E-FABP, psoriasis-associated FABP, PA-FABP), striated muscle and heart (H-FABP, mammary-derived growth inhibitor or MDGI), intestine (I-FABP), liver (L-FABP or FABP1), myelin (M-FABP) and testis (T-FABP). FABP1 (L-FABP) expression is modulated by developmental, hormonal, dietary and pharmacological factors, and is required for cholesterol synthesis and metabolism.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.