

Recombinant Cytokeratin 15 (Esophageal Squamous Cell Carcinoma Marker) Antibody

Rabbit Monoclonal Antibody [Clone KRT15/9088R]

| Catalog No | Format | Size |
|------------------|---|--------|
| 3866-RBM15-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3866-RBM15-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3866-RBM15-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details


| | |
|------------------------|---|
| Clone | KRT15/9088R |
| Gene Name | KRT15 |
| Immunogen | Recombinant human full-length KRT15 protein |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | 52kDa |
| Cellular Localization | Cytoplasm. |
| Species Reactivity | Human |
| Positive Control | A431 cells. Human skin or colon. HeLa |

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant Cytokeratin 15 (Esophageal Squamous Cell Carcinoma Marker) Antibody



Formalin-fixed, paraffin-embedded human prostate cancer stained with Cytokeratin 15 Rabbit Recombinant Monoclonal Antibody (KRT15/9088R). Inset: PBS instead of primary antibody; secondary only negative control.



Formalin-fixed, paraffin-embedded human skin stained with Cytokeratin 15 Rabbit Recombinant Monoclonal Antibody (KRT15/9088R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified KRT15 Rabbit Recombinant Monoclonal Antibody (KRT15/9088R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Keratin 15 is a type I keratin which is expressed only in basal keratinocytes in stratified epithelia and does not appear to have a natural type II expression partner. KRT15 is downregulated in activated keratinocytes. It is a specific marker of stem cells of the hair-follicle bulge and may be a useful marker for diagnosis between basal cell carcinoma (BCC) and trichoepithelioma. Trichoblastoma are benign neoplasms of follicular differentiation frequently found in nevus sebaceous. Many morphologic features are shared with nodular basal cell carcinoma, sometimes rendering a diagnosis difficult. Trichoblastoma and BCC show variable expression of Cytokeratin 15 and Cytokeratin 19, and absence of hair keratins.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Basal Cell Marker