

CD14 (Monocyte / Macrophage Marker) Antibody

Mouse Monoclonal Antibody [Clone LPSR/7738]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 929-MSM16-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 929-MSM16-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 929-MSM16-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |



| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details

| | |
|-------------------------------|--|
| Clone | LPSR/7738 |
| Gene Name | CD14 |
| Immunogen | Recombinant fragment of human CD14 protein (around aa 1-200) (exact sequence is proprietary) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2 / Kappa |
| Mol. Weight of Antigen | 55kDa |
| Cellular Localization | Cell surface |
| Species Reactivity | Human |
| Positive Control | A549 THP-1 cells and human monocytes. Lymph nodes and tonsils. |


*Optimal dilution for a specific application should be determined.

Product Images for CD14 (Monocyte / Macrophage Marker) Antibody



Formalin-fixed, paraffin-embedded human tonsil stained with CD14 Mouse Monoclonal Antibody (LPSR/7738). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

SDS-PAGE Analysis of Purified CD14 Mouse Monoclonal Antibody (LPSR/7738). Confirmation of Integrity and Purity of Antibody.



Formalin-fixed, paraffin-embedded human lymph node stained with CD14 Mouse Monoclonal Antibody (LPSR/7738). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Recognizes a protein of 55kDa, identified as CD14 (also known lipopolysaccharide receptor). CD14 is expressed strongly on monocytes and macrophage and weakly on the surface of neutrophils. CD14 is anchored to cells by linkage to glycosylphosphatidylinositol (GPI) and functions as a high affinity receptor for complexes of LPS and LPS binding protein (LBP). Soluble CD14, also binding to LPS, acts at physiological concentration as an LPS agonist and has, at higher concentrations, an LPS antagonizing effect in cell activation.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

B Cell Markers, Cardiovascular, Dendritic Cell Marker, Hematopoietic Stem Cells, Immunology, Infectious Disease, MAPK Signaling, Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.