

HLA-G (Major Histocompatibility Complex, class I, G) Antibody

Mouse Monoclonal Antibody [Clone HLAG/7749]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 3135-MSM-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3135-MSM9-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3135-MSM9-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|---|
| Clone | HLAG/7749 |
| Gene Name | HLA-G |
| Immunogen | Recombinant fragment (around aa1-200) of human HLAG protein (exact sequence is proprietary) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 39kDa |
| Cellular Localization | Cell membrane, Secreted |
| Species Reactivity | Human |
| Positive Control | Raji cells. Expressed in trophoblasts. Brain. |

*Optimal dilution for a specific application should be determined.


Product Images for HLA-G (Major Histocompatibility Complex, class I, G) Antibody



Formalin-fixed, paraffin-embedded human placenta stained with HLA-G Mouse Monoclonal Antibody (HLAG/7749). Inset: PBS instead of primary antibody; secondary only negative control.



Western blot analysis of Human Brain lysate using HLA-G Mouse Monoclonal Antibody (HLAG/7749).



SDS-PAGE Analysis Purified HLA-G Mouse Monoclonal Antibody (HLA-G/7749).
Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Major histocompatibility complex (MHC), human leukocyte antigen (HLA) molecules are cell-surface receptors that bind foreign peptides and present them to T lymphocytes. MHC class I molecules consist of two polypeptide chains, an α or heavy chain, and a non-covalently associated protein, β 2-microglobulin. Cytotoxic T lymphocytes bind antigenic peptides presented by MHC class I molecules. Antigens that bind to MHC class I molecules are typically 8-10 residues in length and are stabilized in a peptide binding groove. MHC class II molecules are encoded by polymorphic MHC genes and consist of a noncovalent complex of an α and β chain. Helper T lymphocytes bind antigenic peptides presented by MHC class II molecules. MHC class II molecules bind 13-18 amino acid antigenic peptides. Accumulating in endosomal/lysosomal compartments and on the surface of B cells, HLA-DM and -DO molecules regulate binding of exogenous peptides to class II molecules (HLA-DR) by sustaining a conformation that favors peptide exchange. The differential structural properties of MHC class I and class II molecules account for their respective roles in activating different populations of T lymphocytes.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Cytokine Signaling, Immunology, Infectious Disease