

Recombinant Bax (Apoptosis Marker) Antibody

Rabbit Monoclonal Antibody [Clone BAX/3158R]

Catalog No	Format	Size
581-RBM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20ug
581-RBM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100ug
581-RBM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100ug


Applications	Tested Dillution	Note
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	BAX/3158R
Gene Name	BAX
Immunogen	Recombinant fragment (around aa1-100) of human BAX protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	21kDa
Cellular Localization	Cytoplasm
Species Reactivity	Human
Positive Control	A549, MCF7, HL-60, Jurkat, K562 or HeLa cells. Reed-Sternberg cells in Hodgkin's lymphoma, Human Intestine, Human Lung, Human Kidney, Human Spleen, A549, MCF7

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant Bax (Apoptosis Marker) Antibody




SDS-PAGE Analysis of Purified BAX Recombinant Rabbit Monoclonal Antibody (BAX/3158R). Confirmation of Purity and Integrity of Antibody



Western Blot Analysis of A549 and MCF7 cell lysates using Bax (Apoptosis Marker) Recombinant Rabbit Monoclonal Antibody (BAX/3158R).



Western Blot Analysis of Human Intestine, Human Lung, Human Kidney, Human Spleen, Human Stomach, A549 and MCF7 lysates using BAX Recombinant Rabbit Monoclonal Antibody (BAX/3158R).



Western Blot Analysis of Human Intestine, Mouse Intestine, Rat Intestine, Hamster Intestine, Guinea Pig Intestine, Rabbit Intestine, Human Kidney, Mouse Kidney, Rat Kidney, Hamster Kidney, Guinea pig Kidney, Rabbit Kidney, Human Spleen, Mouse Spleen, Rat Spleen, Hamster Spleen, Guinea pig Spleen and Rabbit Spleen tissue lysates using BAX Recombinant Rabbit Monoclonal Antibody (BAX/3158R).

Specificity & Comments

Recognizes a protein of 21 kDa, identified as the Bax protein. This MAb is highly specific to Bax and shows no cross-reaction with Bcl-2 or Bcl-X protein. Bcl-2 blocks cell death following a variety of stimuli. Bax has extensive amino acid homology with Bcl-2 and it homodimerizes and forms heterodimers with Bcl-2. Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3 dependent cell line, and Bax also counters the death repressor activity of Bcl-2.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Apoptosis, Autophagy, Breast Cancer, Cardiovascular, Colon Cancer, Complement System, Lung Cancer, Ovarian Cancer, Signal Transduction, Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.