Nucleolin (Marker of Human Cells) Antibody

Mouse Monoclonal Antibody [Clone NCL/902]

Catalog No	Format	Size
4691-MSM1-CF488-100T	Purified Ab conjugated to CF488	0.5 ml at 100ug/ml


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	NCL/902
Gene Name	NCL
Immunogen	Recombinant full-length human NCL protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	76kDa
Cellular Localization	Cytoplasm, nucleoli, Nucleolus, Nucleus
Species Reactivity	Human
Positive Control	HeLa cells. Breast Cancer.

**Optimal dilution for a specific application should be determined.*


Product Images for Nucleolin (Marker of Human Cells) Antibody



Flow Cytometric Analysis of human Nucleolin on 293T cells. Black: cells alone; Gray: Isotype Control; Green: CF488-labeled Nucleolin Monoclonal Antibody (NCL/902).



Formalin-fixed, paraffin-embedded human colon carcinoma stained with CF488-labeled Nucleolin Monoclonal Antibody (NCL/902).



Flow cytometric analysis of human Nucleolin antigen on Jurkat cells. Grey: isotype control; green: FITC-labeled; purple: APC-labeled; Turquoise: AF700-labeled Nucleolin Mouse Monoclonal Antibody (NCL/902).

Specificity & Comments

Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleolar chromatin and pre-ribosomal particles. Human NCL gene consists of 14 exons with 13 introns and spans approximately 11kb. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly.This Mouse Monoclonal Antibody can be used to stain the nucleoli in cell or tissue preparations and can be used as a marker of the nucleoli in subcellular fractions. It produces a speckled pattern in the nuclei of cells of normal and malignant cells and may be used to stain the nucleoli of cells in fixed or frozen tissue sections. It can be used with paraformaldehyde fixed frozen tissue or cell preparations and formalin fixed, paraffin-embedded tissue sections.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Nuclear Marker

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.