

CD86 (Dendritic Cells Maturation Marker) Antibody

Mouse Monoclonal Antibody [Clone BU63]

| Catalog No | Format | Size |
|------------------|------------------------------|--------------------|
| 942-MSM1-PE-100T | Purified Ab Conjugated to PE | 0.5 ml at 100ug/ml |


| Applications | Tested Dillution | Note |
|-------------------------|---------------------|------|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |

Product Details


| | |
|-------------------------------|------------------------------------------------------|
| Clone | BU63 |
| Gene Name | CD86 |
| Immunogen | ARH-77 (B-lymphoblastoid cell line) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 70kDa |
| Cellular Localization | Cell membrane, Cell surface |
| Species Reactivity | Human |
| Positive Control | 293T or Jurkat cells. Monocytes and dendritic cells. |

*Optimal dilution for a specific application should be determined.

Product Images for CD86 (Dendritic Cells Maturation Marker) Antibody



SDS-PAGE Analysis of Purified CD86 Mouse Monoclonal Antibody (BU63). Confirmation of Integrity and Purity of Antibody.



Flow cytometric analysis of human peripheral blood mononuclear cell (PBMCs). Red: PE-labeled CD86 Mouse Monoclonal Antibody (BU63); Green: Isotype Control.

Specificity & Comments

Recognizes a protein of 70kDa, which is identified as CD86 (HLDA V; WS Code BP BP072. HLDA V; WS Code A A109. HLDA VI; WS Code BP 95. HLDA VI; WS Code B CD86.9). CD86 is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. CD86 expression is rapidly upregulated by B cell specific stimuli with peak expression at 18 to 42 hours after stimulation. CD86, along with CD80/B71, is an important accessory molecule in T cell co-stimulation via its interaction with CD28 and CD152/CTLA4. Since CD86 has rapid kinetics of induction, it is believed to be the major CD28 ligand expressed early in the immune response. It is also found on malignant Hodgkin and Reed Sternberg (HRS) cells in Hodgkin's disease.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

B Cell Markers, Cardiovascular, Cytokine Signaling, Dendritic Cell Marker, Hematopoietic Stem Cells, Immunology, Infectious Disease, Signal Transduction
