

Periostin (POSTN) Antibody

Mouse Monoclonal Antibody [Clone POSTN/4095]

Catalog No	Format	Size
10631-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10631-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10631-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	POSTN/4095
Gene Name	POSTN
Immunogen	Recombinant fragment (around aa193-326) of human POSTN protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	84/74kDa (Periostin); 90kDa (secreted glycoprotein)
Cellular Localization	Extracellular matrix, Extracellular space, Secreted
Species Reactivity	Human
Positive Control	Human placenta, breast or colon tissue.

*Optimal dilution for a specific application should be determined.

Product Images for Periostin (POSTN) Antibody



Formalin-fixed, paraffin-embedded human adrenal gland stained with Periostin Mouse Monoclonal Antibody (POSTN/4095). HIER: Tris/EDTA, pH9.0, 45min. 2°: HRP-polymer, 30min. DAB, 5min.

SDS-PAGE Analysis Purified Periostin Mouse Monoclonal Antibody (POSTN/4095). Confirmation of Purity and Integrity of Antibody

Specificity & Comments

Periostin (PN), also designated osteoblast-specific factor 2 (OSF-2), is a disulfide linked protein originally isolated as a osteoblast-specific factor. Periostin is a secreted protein that binds heparin and functions as a ligand for $\alpha 3$ and $\alpha 5$ integrins. In preosteoblasts, Periostin acts as a cell adhesion molecule and plays a role in osteoblast recruitment, spreading and attachment. Periostin is mainly detected in lower gastrointestinal tract, aorta, stomach, placenta, uterus and breast tissues but is up-regulated in epithelial ovarian tumors and overexpressed in breast cancer. Expression of Periostin is increased by bone morphogenetic protein (BMP2) and transforming growth factor $\beta 1$ (TGF $\beta 1$). Periostin contains a typical signal sequence, followed by a cysteine-rich domain, a fourfold repeated domain, which shows homology with the insect protein fasciclin, and a C-terminal domain.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Endothelial Cell Marker
