

Recombinant P-Cadherin (CDH3) (Basal Epithelial Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone r56C1]

Catalog No	Format	Size
1001-MSM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1001-MSM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1001-MSM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	r56C1
Gene Name	CDH3
Immunogen	Prokaryotic recombinant protein corresponding to a region of the external domain of the P-cadherin molecule
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	116kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	A431 or PC-3 cells. Human pancreas, placenta or prostate.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant P-Cadherin (CDH3) (Basal Epithelial Cell Marker) Antibody



SDS-PAGE Analysis of Purified P-Cadherin Recombinant Mouse Monoclonal Antibody (r56C1). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human colon carcinoma stained with P-Cadherin Recombinant Mouse Monoclonal Antibody (r56C1). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Recognizes a protein of 116kDa, identified as P-Cadherin-1 (CDH3). It is a calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a six-cadherin cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene have been associated with congenital hypotrichosis with juvenile macular dystrophy.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Basal Cell Marker
