

ABCC4 / ATP Binding Cassette Subfamily C Member 4 (Drug Target) Antibody

Mouse Monoclonal Antibody [Clone ABCC4/9180]

Catalog No	Format	Size
10257-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10257-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10257-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
--------------	------------------	------

Product Details


Clone	ABCC4/9180
Gene Name	ABCC4
Immunogen	Recombinant fragment (around aa 1062-1174) of human ABCC4 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	150kDa
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	Human prostate. Kidney.

**Optimal dilution for a specific application should be determined.*


Product Images for ABCC4 / ATP Binding Cassette Subfamily C Member 4 (Drug Target) Antibody




Formalin-fixed, paraffin-embedded human prostate stained with ABCC4 Mouse Monoclonal Antibody (ABCC4/9180). HIER: Tris/EDTA, pH9.0, 45min. 2°: HRP-polymer, 30min. DAB, 5min.



Western blot analysis of human kidney tissue lysates using ABCC4 Mouse Monoclonal Antibody (ABCC4/9180).



Western blot analysis of kidney tissue lysates from human, rat and rabbit using ABCC4 Mouse Monoclonal Antibody (ABCC4/9180).



SDS-PAGE Analysis of Purified ABCC4 Mouse Monoclonal Antibody (ABCC4/9180). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

The two members of the large family of ABC transporters known to confer multidrug resistance in human cancer cells are the Mdr-1 P-glycoprotein and the multidrug-resistance protein MRP1. MRP1 is an integral membrane protein that contains an MDR-like core, an N-terminal membrane-bound region and a cytoplasmic linker, and it is expressed in various cerebral cells, as well as in lung, testis and peripheral blood. The MRP gene family also includes MRP2, which is alternatively designated cMOAT (for canalicular multispecific organic anion transporter), and MRP3, which are both conjugate export pumps expressed predominantly in hepatocytes. MRP2 localizes exclusively to the apical membrane and is constitutively expressed at a high level in normal liver cells. Conversely, MRP3 localizes to the basolateral membrane where it also mediates the transport of the organic anion S-(2,4-dinitrophenyl-) glutathione toward the basolateral side of the membrane. MRP3 is normally expressed at comparatively lower levels than MRP2 and increases only when secretion across the apical membrane by MRP2 is impaired. MRP6 is highly expressed in liver and kidney, whereas MRP4 and MRP5 are detected in various tissues, yet at much lower levels of expression.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular