

## DBC2 / RHOBTB2 (Tumor Suppressor) Antibody

Mouse Monoclonal Antibody [Clone DBC2/12581]

Catalog No	Format	Size
23221-MSM16-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
23221-MSM16-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
23221-MSM16-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note

### Product Details


<b>Clone</b>	DBC2/12581
<b>Gene Name</b>	RHOBTB2
<b>Immunogen</b>	Human recombinant RHOBTB2 protein fragment (around aa500-700) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	83kDa
<b>Cellular Localization</b>	Cytoplasm, Secreted in plasma
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Human heart muscle or brain tissue.

\*Optimal dilution for a specific application should be determined.

### Product Images for DBC2 / RHOBTB2 (Tumor Suppressor) Antibody



SDS-PAGE Analysis of Purified Rho-related BTB domain-containing protein 2 Mouse Monoclonal Antibody (DBC2/12581). Confirmation of Purity and Integrity of Antibody.



SDS-PAGE Analysis of Purified RHOBTB2 Mouse Monoclonal Antibody (DBC2/12581). Confirmation of Purity and Integrity of Antibody.

### **Specificity & Comments**

The Rho subfamily of Ras-related GTPases controls multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. DBC-2 (deleted in breast cancer 2 gene protein), also known as RHOBTB2 (Rho-related BTB domain-containing protein 2), is a 727 amino acid member of the RhoBTB subfamily of Rho GTPases. Members of the RhoBTB subfamily are evolutionarily conserved and are characterized by a proline-rich region, a GTPase domain and two tandem BTB repeats. Expressed ubiquitously with highest levels in neural tissue, heart, brain and fetal lung, DBC-2 contains two BTB (POZ) domains through which it may bind to and regulate the function of target proteins, such as CUL-3. Additionally, DBC-2 is thought to function as a regulator of cell cycle and apoptosis events. Under normal conditions, DBC-2 is thought to exhibit tumor suppressor activity. Mutations in the gene encoding DBC-2 are associated with breast cancer, suggesting that mutated DBC-2 may play a role in carcinogenesis.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### **Research Areas**

Signal Transduction

---