

Recombinant Glutamine Synthetase / GLUL Antibody

Rabbit Monoclonal Antibody [Clone GLUL/12902R]

Catalog No	Format	Size
2752-RBM39-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2752-RBM39-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2752-RBM39-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	GLUL/12902R
Gene Name	GLUL
Immunogen	Recombinant fragment (around aa 100-200) of human GLUL protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	42kDa
Cellular Localization	Cytoplasm, Mitochondrion
Species Reactivity	Human
Positive Control	Y79 cells. Human brain, placenta, testis or stomach.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Glutamine Synthetase / GLUL Antibody



Formalin-fixed, paraffin-embedded human stomach stained with Glutamine Synthetase Recombinant Rabbit Monoclonal Antibody (GLUL/12902R). Inset: PBS instead of primary antibody; secondary only negative control.

Western blot analysis of human brain tissue lysate using Glutamine Synthetase Recombinant Rabbit Monoclonal Antibody (GLUL/12902R).

Specificity & Comments

Glutamine synthetase (Gl Syn) forms a homo-octamer that serves as a catalyst for the amination of glutamic acid to form glutamine. This enzyme is a marker for astrocytes, which serve as the primary site of conversion of glutamic acid to glutamine in the brain. Induction of glutamine synthetase is seen upon astrocyte cell contact with neurons. Elevated expression of glutamine synthetase in glial cells has been shown to protect neurons from degeneration due to excess glutamate. Glutamine synthetase is also present in the liver and is involved in nitrogen homeostasis. Overexpression of glutamine synthetase has been shown in primary liver cancers, indicating a potential role for glutamine synthetase in hepatocyte transformation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Mitochondria Marker, Neuroinflammation
