

Hepatocyte nuclear factor 1 beta (HNF1b) / Transcription Factor 2 (TCF2) Antibody

Mouse Monoclonal Antibody [Clone HNF1B/9518]

Catalog No	Format	Size
6928-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6928-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6928-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	HNF1B/9518
Gene Name	HNF1B
Immunogen	Recombinant fragment (around aa1-200) of human HNF1B protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	61kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Human kidney, pancreas or liver tissue.

**Optimal dilution for a specific application should be determined.*

Product Images for Hepatocyte nuclear factor 1 beta (HNF1b) / Transcription Factor 2 (TCF2) Antibody



SDS-PAGE Analysis of Purified HNF1B Mouse Monoclonal Antibody (HNF1B/9519). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human liver carcinoma stained with HNF1B Mouse Monoclonal Antibody (HNF1B/9519). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

HNF-1 (a and b), HNF-3 (a, b and g), HNF-4 (a and g) and HNF-6 compose, in part, a homeoprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3a, HNF-3b and HNF-3g regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3a and HNF-3b have also been shown to be involved in gastrulation events such as body axis formation. HNF-4a and HNF-4g have been shown to be important for early embryo development. HNF-4a is expressed in liver, kidney, pancreas, small intestine, testis and colon; and HNF-4g is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3b, which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transcriptional activator for at least 22 other hepatocyte enriched genes, including cytochrome P450 2C13 and a-1 antitrypsin.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Kidney, Pancreatic Cancer, Transcription Factors
