

## FABP2 (Marker of Metastatic Potential in Colorectal Cancer) Antibody

Mouse Monoclonal Antibody [Clone FABP2/6344]

Catalog No	Format	Size
2169-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2169-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2169-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	FABP2/6344
<b>Gene Name</b>	FABP2
<b>Immunogen</b>	Recombinant fragment (around aa1-132) of human FABP2 protein (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2 / Kappa
<b>Mol. Weight of Antigen</b>	15.21kDa
<b>Cellular Localization</b>	Cytoplasm
<b>Species Reactivity</b>	Guinea Pig, Hamster, Human, Mouse, Rabbit, Rat
<b>Positive Control</b>	MOLT-4 or HEK-293 cells. esophagus or colon. Human Intestine

\*Optimal dilution for a specific application should be determined.



### Product Images for FABP2 (Marker of Metastatic Potential in Colorectal Cancer) Antibody



Formalin-fixed, paraffin-embedded human small intestine stained with FABP2 Mouse Monoclonal Antibody (FABP2/6344). HIER: Tris/EDTA, pH9.0, 45min. 2°: HRP-polymer, 30min. DAB, 5min.




SDS-PAGE Analysis of Purified FABP2 Mouse Monoclonal Antibody (FABP2/6344). Confirmation of Purity and Integrity of Antibody.




Analysis of Protein Array containing more than 19,000 full-length human proteins using FABP2 Mouse Monoclonal Antibody (FABP2/6344). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Western blot analysis of Human Intestine, Mouse Intestine and Rat Intestine tissue lysates using FABP2 Mouse Monoclonal Antibody (FABP2/6344).



Western blot analysis of Human intestine, Mouse intestine, Rat intestine, Hamster intestine, Guinea pig intestine and Rabbit intestine tissue lysates using FABP2 Mouse Monoclonal Antibody (FABP2/6344).



Western blot analysis of human small intestine tissue lysate using FABP2 Mouse Monoclonal Antibody (FABP2/6344).

## Specificity & Comments

The intracellular fatty acid-binding proteins (FABPs) belong to a multigene family with nearly twenty identified members. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Intestinal fatty acid-binding protein 2 gene contains four exons and is an abundant cytosolic protein in small intestine epithelial cells. This gene has a polymorphism at codon 54 that identified an alanine-encoding allele and a threonine-encoding allele. Thr-54 protein is associated with increased fat oxidation and insulin resistance.

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

## Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

## Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## Research Areas

Metabolism