

## Recombinant Folate Receptor Alpha (FRalpha) / FOLR1 Antibody


Mouse Monoclonal Antibody [Clone rFOLR1/13324]

Catalog No	Format	Size
2348-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2348-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2348-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
<b>Product Details</b>		
Clone	rFOLR1/13324	
Gene Name	FOLR1	
Immunogen	Recombinant human FOLR1 protein	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG2a / Kappa	
Mol. Weight of Antigen	29kDa	
Cellular Localization	Cell membrane, Cytoplasm, Secreted	
Species Reactivity	Human	
Positive Control	Human kidney, thymus, placenta or lung.	

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant Folate Receptor Alpha (FRalpha) / FOLR1 Antibody



SDS-PAGE Analysis of Purified Folate Receptor Alpha Recombinant Mouse Monoclonal Antibody (rFOLR1/13324). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

Folate receptor alpha (FR?) is a membrane-bound protein that facilitates high-affinity folate transport, essential for cell metabolism, DNA synthesis, and repair, particularly in rapidly dividing cancer cells with increased folate demands. This receptor, encoded by the FOLR1 gene, employs a cotransport mechanism that leads to both membranous and cytoplasmic localization and is selectively overexpressed in certain epithelial malignancies while remaining highly restricted in normal tissues. FR? expression correlates with tumor stage and grade, suggesting a role in tumor progression by enhancing folate uptake or generating regulatory signals. Its regulation is influenced by extracellular folate levels, homocysteine accumulation, hormones, transcription factors, and genetic mutations. Members of the folate receptor family share highly conserved sequences in the open reading frames but differ in amino acids in the 5' untranslated regions and as a consequence can differ in function and tissue expression. Given the selective overexpression of FR? in tumors and its association with aggressive cancer phenotypes, FR? is a valuable marker for studying tumor biology and progression, particularly in epithelial cancers.

### Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Immuno Oncology

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---