

Recombinant MCM5 (Transcription Factor) Antibody

Mouse Monoclonal Antibody [Clone rMCM5/13646]

Catalog No	Format	Size
4174-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4174-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4174-MSM4-P1ABX	Purified Ab without BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	



Product Details

Clone	rMCM5/13646
Immunogen	Recombinant fragment (around aa500-734) of human MCM5 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	90kDa
Cellular Localization	Chromosome, Nucleus
Species Reactivity	Human
Positive Control	A-431 cells. Human tonsil, spleen or lymph node.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant MCM5 (Transcription Factor) Antibody

Purity: SDS-PAGE



SDS-PAGE Analysis of Purified MCM5 Recombinant Mouse Monoclonal Antibody (rMCM5/13646). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human colon stained with MCM5 Recombinant Mouse Monoclonal Antibody (rMCM5/13646). Inset: PBS instead of primary antibody; secondary only negative control.



Western Blot Analysis of A-431 cell lysate using MCM5 Recombinant Mouse Monoclonal Antibody (rMCM5/13646).

Specificity & Comments

The mini-chromosome maintenance (MCM) family of proteins, including MCM2, MCM3, MCM4 (Cdc21), MCM5 (Cdc46), MCM6 (Mis5) and MCM7 (Cdc47), are regulators of DNA replication that act to ensure replication occurs only once in the cell cycle. Expression of MCM proteins increases during cell growth, peaking at G1 to S phase. The MCM proteins each contain an ATP-binding motif, which is predicted to mediate ATP-dependent opening of double-stranded DNA. MCM proteins are regulated by E2F transcription factors, which induce MCM expression, and by protein kinases, which interact with MCM proteins to maintain the postreplicative state of the cell. MCM2/MCM4 complexes function as substrates for Cdc2/cyclin B in vitro. Cleavage of MCM3, which can be prevented by caspase inhibitors, results in the inactivation of the MCM complex (composed of at least MCM proteins 2-6) during apoptosis. A complex composed of MCM4, MCM6 and MCM7 has been shown to be involved in DNA helicase activity; and MCM5 is involved in IFN- γ -induced Stat1 transcription activation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Transcription Factors