

## Recombinant CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody

Rabbit Monoclonal Antibody [Clone CD70/12949R]

Catalog No	Format	Size
970-RBM26-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
970-RBM26-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
970-RBM26-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	CD70/12949R
<b>Immunogen</b>	Recombinant fragment (around aa176-193) of the human CD70 protein (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	21.12kDa
<b>Cellular Localization</b>	Cell membrane
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Raji cells. Lymph node or tonsil. HeLa or Jurkat.

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody



Western Blot Analysis of HeLa and Jurkat cell lysates using CD70 Recombinant Rabbit Monoclonal Antibody (CD70/12949R).



SDS-PAGE Analysis of Purified CD70 Recombinant Rabbit Monoclonal Antibody (CD70/12949R). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

It recognizes a protein of 30kDa, identified as CD70. It is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T- and B-lymphocytes. It induces proliferation of co-stimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.