

AIF1 / Iba1 (Microglia Marker) Antibody

Mouse Monoclonal Antibody [Clone MSVA-955M]

Catalog No	Format	Size
199-MSM19-P0	Purified Ab with BSA and Azide	20 ug
199-MSM19-P1	Purified Ab with BSA and Azide	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1:100-1:200	Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.

Product Details	
Clone	MSVA-955M
Immunogen	Purified recombinant human AIF1 protein (around aa 1-146) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2c / Kappa
Mol. Weight of Antigen	17kDa
Cellular Localization	Cell projection, Cytoplasm, Cytoskeleton, Phagocytic cup, Ruffle membrane
Species Reactivity	Human
Positive Control	Colon: An at least moderate to strong cytoplasmic AIF1 immunostaining should be seen in mucosal macrophages.

*Optimal dilution for a specific application should be determined.

Product Images for AIF1 / Iba1 (Microglia Marker) Antibody



Allograft inflammatory factor 1 Mouse Monoclonal Antibody (MSVA-955M) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

Specificity & Comments

AIF1 is a cytoplasmic, calcium-binding protein that is thought to play a role in macrophage activation and function. AIF1, containing two EF domains, is induced by cytokines and Interferons. In an unstimulated state, AIF1 colocalizes with actin, and upon stimulation, translocates to lamellipodia. It is also a marker of human microglia and is expressed by macrophages in injured skeletal muscle. The gene encoding AIF1 resides in the tumor necrosis factor (TNF) cluster of genes, located in the region represented by the human major histocompatibility complex (MHC).

Supplied As

Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Dendritic Cell Marker

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
