

## CD23 (Fc Epsilon RII) Antibody

Mouse Monoclonal Antibody [Clone MSVA-023M]

| Catalog No    | Format                         | Size   |
|---------------|--------------------------------|--------|
| 2208-MSM26-P0 | Purified Ab with BSA and Azide | 20 ug  |
| 2208-MSM26-P1 | Purified Ab with BSA and Azide | 100 ug |


| Applications               | Tested Dillution | Note   |
|----------------------------|------------------|--|
| Immunohistochemistry (IHC) | 1:100-1:200      | Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions. |

### Product Details

|                               |  |
|-------------------------------|--|
| <b>Clone</b>                  | MSVA-023M  |
| <b>Immunogen</b>              | Recombinant fragment (around aa 48-321) of human FCER2/CD23 protein (exact sequence is proprietary)  |
| <b>Host</b>                   | Mouse  |
| <b>Clonality</b>              | Monoclonal   |
| <b>Isotype / Light Chain</b>  | IgG1 / Kappa   |
| <b>Mol. Weight of Antigen</b> | 37kDa (soluble form); 45kDa (membrane-bound form)  |
| <b>Cellular Localization</b>  | Cell membrane, Secreted  |
| <b>Species Reactivity</b>     | Human  |
| <b>Positive Control</b>       | Tonsil: The follicular dendritic cells of the germinal centres should show a strong CD23 immunostaining while the majority of lymphocytes in the mantle zone of the follicles should show an at least weak staining. |

\*Optimal dilution for a specific application should be determined.

### Product Images for CD23 (Fc Epsilon RII) Antibody



Low affinity immunoglobulin epsilon Fc receptor Mouse Monoclonal Antibody (MSVA-023M) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

**Specificity & Comments**

CD23 (FCE2) is a type II integral membrane glycoprotein that is expressed on mature B cells, monocytes, eosinophils, platelets and dendritic cells. CD23 is a low affinity IgE receptor that mediates IgE-dependent cytotoxicity and phagocytosis by macrophages and eosinophils. CD23 associates as an oligomer where cooperative binding of at least two lectin domains is required for high affinity IgE binding to CD23. It may play a role in antigen presentation by B cells by interacting with CD40. CD23 has been shown to be associated with the Fyn tyrosine kinase. The truncated molecule can be secreted, then function as a potent mitogenic growth factor. CD23 is expressed on a subpopulation of peripheral blood cells, B-lymphocytes and on EBV transformed B lymphoblastoid cell lines. CD23 is also detected in neoplastic cells from cases of B cell chronic lymphocytic leukemia and some cases on centroblastic/centrocytic lymphoma.

---

**Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

**Supplied As**

Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

---

**Storage and Stability**

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

**Research Areas**

B Cell Markers, Complement System, Cytokine Signaling, Immunology, Mast Cell Marker, Signal Transduction

---