

Desmoglein-3 (Squamous Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone MSVA-543M]

Catalog No	Format	Size
1830-MSM17-P0	Purified Ab with BSA and Azide	20 ug
1830-MSM17-P1	Purified Ab with BSA and Azide	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1:75-1:150	Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:100 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.

Product Details

Clone	MSVA-543M
Immunogen	Recombinant fragment (around aa 379-491) human DSG3 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	130kDa
Cellular Localization	Cell junction, Cell membrane, Desmosome
Species Reactivity	Human
Positive Control	Tonsil: A moderate to strong desmoglein 3 immunostaining should be seen in a fraction of squamous epithelial cells.

*Optimal dilution for a specific application should be determined.

Product Images for Desmoglein-3 (Squamous Cell Marker) Antibody



Desmoglein-3 Mouse Monoclonal Antibody (MSVA-543M) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

Specificity & Comments

Recognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.

Supplied As

Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
