

Recombinant Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone MSVA-561R]


Catalog No	Format	Size
3623-RBM15-P0	Purified Ab with BSA and Azide	20 ug
3623-RBM15-P1	Purified Ab with BSA and Azide	100 ug

Applications	Tested Dilution	Note
Immunohistochemistry (IHC)	1:75-1:150	Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:50 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.

Product Details	
Clone	MSVA-561R
Immunogen	Recombinant fragment of human Inhibin alpha protein (around aa 233-362) (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	47kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Adrenal gland: A moderate to strong INHA positivity should be seen in adrenocortical cells.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody



Inhibin alpha chain Rabbit Recombinant Monoclonal Antibody (MSVA-561R) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

Specificity & Comments

It recognizes a 47kDa protein, which is identified as alpha sub-unit of Inhibin. It is a gonadal protein that preferentially suppresses the secretion of pituitary follicle-stimulating hormone (FSH). Inhibin comprises two subunits, Inhibin A and Inhibin B. Each subunit consists of the same ? subunit, covalently linked to 1 of 2 distinct subunits, ?? or ??. Originally isolated from ovarian follicular fluid and characterized as a disulphide-linked dimeric glycoprotein, inhibin belongs to the transforming growth factor ? (TFG?) superfamily. Antibodies against Inhibin are useful in making a differentiation between adrenal cortical tumors and renal cell carcinoma. Sex cord stromal tumors of the ovary as well as trophoblastic tumors also demonstrate cytoplasmic positivity. Inhibin antibody is also used to make the differential diagnosis of intra-uterine vs. ectopic pregnancy in endometrial curetting.

Supplied As

Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
