

## Recombinant PGP9.5 / UchL1 (pan-Neuronal Marker) Antibody

Rabbit Monoclonal Antibody [Clone MSVA-905R]

| Catalog No    | Format                         | Size   |
|---------------|--------------------------------|--------|
| 7345-RBM52-P0 | Purified Ab with BSA and Azide | 20 ug  |
| 7345-RBM52-P1 | Purified Ab with BSA and Azide | 100 ug |

| Applications               | Tested Dillution | Note                                                                                                                                                                                                                                                                                                                                                                                                       |
|----------------------------|------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunohistochemistry (IHC) | 1:100-1:200      | Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions. |

### Product Details

|                               |                                                                                          |
|-------------------------------|------------------------------------------------------------------------------------------|
| <b>Clone</b>                  | MSVA-905R                                                                                |
| <b>Immunogen</b>              | Recombinant full-length human UCHL1 protein                                              |
| <b>Host</b>                   | Rabbit                                                                                   |
| <b>Clonality</b>              | Monoclonal                                                                               |
| <b>Isotype / Light Chain</b>  | IgG / Kappa                                                                              |
| <b>Mol. Weight of Antigen</b> | 20-30kDa                                                                                 |
| <b>Cellular Localization</b>  | Cytoplasm, Endoplasmic reticulum membrane                                                |
| <b>Species Reactivity</b>     | Human                                                                                    |
| <b>Positive Control</b>       | Pancreas: Islets cells and interspersed nerve fibres should show strong PGP9.5 staining. |

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant PGP9.5 / UchL1 (pan-Neuronal Marker) Antibody



Ubiquitin carboxyl-terminal hydrolase isozyme L1 Rabbit Recombinant Monoclonal Antibody (MSVA-905R) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

### Specificity & Comments

This MAbs reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in distal renal tubular epithelium, spermatogonia, Leydig cells, oocytes, melanocytes, prostatic secretory epithelium, ejaculatory duct cells, epididymis, mammary epithelial cells, Merkel cells, and dermal fibroblasts. Furthermore, immunostaining for PGP9.5 has been shown in a wide variety of mesenchymal neoplasms as well. A mutation in PGP9.5 gene is believed to cause a form of Parkinson's disease.

### Supplied As

Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Neuroscience

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---