

## Recombinant S100B (Astrocyte and Melanoma Marker) Antibody

Rabbit Monoclonal Antibody [Clone MSVA-490R]

| Catalog No    | Format                         | Size   |
|---------------|--------------------------------|--------|
| 6285-RBM43-P0 | Purified Ab with BSA and Azide | 20 ug  |
| 6285-RBM43-P1 | Purified Ab with BSA and Azide | 100 ug |


| Applications               | Tested Dillution | Note   |
|----------------------------|------------------|--|
| Immunohistochemistry (IHC) | 1:100-1:200      | Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions. |

### Product Details

|                               |   |
|-------------------------------|---|
| <b>Clone</b>                  | MSVA-490R   |
| <b>Immunogen</b>              | Recombinant full-length human S100B protein   |
| <b>Host</b>                   | Rabbit  |
| <b>Clonality</b>              | Monoclonal  |
| <b>Isotype / Light Chain</b>  | IgG / Kappa   |
| <b>Mol. Weight of Antigen</b> | 10-12kDa  |
| <b>Cellular Localization</b>  | Cytoplasm, Nucleus  |
| <b>Species Reactivity</b>     | Human   |
| <b>Positive Control</b>       | Appendix: Schwann cells of peripheral nerves and adipocytes should show a strong, predominantly cytoplasmic S100B staining. |

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant S100B (Astrocyte and Melanoma Marker) Antibody



Protein S100-B Rabbit Recombinant Monoclonal Antibody (MSVA-490R) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

### Specificity & Comments

S100 belongs to the family of calcium binding proteins. S100A and S100B proteins are two members of the S100 family. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the beta-chain (i.e. in S-100A and S-100B) but not on the alpha-chain of S-100 (i.e. in S-100A and S100A0). This antibody can be used to localize S-100A and S-100B in various tissue sections. S-100 protein has been found in normal melanocytes, Langerhans cells, histiocytes, chondrocytes, lipocytes, skeletal and cardiac muscle, Schwann cells, epithelial and myoepithelial cells of the breast, salivary and sweat glands, as well as in glial cells. Neoplasms derived from these cells also express S-100 protein, albeit non-uniformly. A large number of well-differentiated tumors of the salivary gland, adipose and cartilaginous tissue, and Schwann cell-derived tumors express S-100 protein. Almost all malignant melanomas and cases of histiocytosis X are positive for S-100 protein.

---

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### Supplied As

Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

---

### Storage and Stability

Antibody with azide store at 2 to 8 °C. Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### Research Areas

Cardiovascular, Cytokine Signaling, Dendritic Cell Marker, Immunology, Neural Stem Cells, Neuroscience, Nuclear Marker, Signal Transduction

---