

Uroplakin 1A (Urothelial Differentiation Marker) Antibody

Mouse Monoclonal Antibody [Clone MSVA-735M]

Catalog No	Format	Size
11045-MSM14-P0	Purified Ab with BSA and Azide	20 ug
11045-MSM14-P1	Purified Ab with BSA and Azide	100 ug


Applications	Tested Dilution	Note
Immunohistochemistry (IHC)	1:100-1:200	Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37°C for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.

Product Details

Clone	MSVA-735M
Immunogen	Recombinant fragment (around aa 114-173) of human Uroplakin 1A (UPK1A) protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	29kDa
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	Urinary bladder: A strong membranous and cytoplasmic Upk1a immunostaining should be seen in the urothelium (the staining can be limited to the top 1/3 or 2/3 of the urothelium).

*Optimal dilution for a specific application should be determined.

Product Images for Uroplakin 1A (Urothelial Differentiation Marker) Antibody



Uroplakin-1a Mouse Monoclonal Antibody (MSVA-735M) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

Specificity & Comments

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is found in the asymmetrical unit membrane (AUM) where it can complex with other transmembrane 4 superfamily proteins. It may play a role in normal bladder epithelial physiology, possibly in regulating membrane permeability of superficial umbrella cells or in stabilizing the apical membrane through AUM/cytoskeletal interactions. The protein may also play a role in tumor suppression.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
