

Nectin 4 / Nectin Cell Adhesion Molecule 4 / PVRL4 Antibody

Mouse Monoclonal Antibody [Clone NECTIN4/9906]

Catalog No	Format	Size
81607-MSM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
81607-MSM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
81607-MSM6-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	NECTIN4/9906
Gene Name	NECTIN4
Immunogen	Recombinant fragment (around aa1-200) of human NECTIN4 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	66kDa
Cellular Localization	Adherens junction, Cell junction, Cell membrane, Secreted
Species Reactivity	Human
Positive Control	Human tonsil, placenta or breast. A-431, MCF7

*Optimal dilution for a specific application should be determined.


Product Images for Nectin 4 / Nectin Cell Adhesion Molecule 4 / PVRL4 Antibody




Formalin-fixed, paraffin-embedded human skin stained with Nectin 4 Mouse Monoclonal Antibody (NECTIN4/9906). Inset: PBS instead of primary antibody; secondary only negative control.




Western blot analysis of A-431 and MCF7 cell lysates using NECTIN 4 Mouse Monoclonal Antibody (NECTIN4/9906).



SDS-PAGE Analysis of Purified Nectin 4 Mouse Monoclonal Antibody (NECTIN4/9906). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using Nectin 4 Mouse Monoclonal Antibody (NECTIN4/9906). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human placenta stained with Nectin 4 Mouse Monoclonal Antibody (NECTIN4/9906). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Homologous to the poliovirus receptor (PVR/CD155), the nectin immunoglobulin superfamily comprises four known isoforms (-1, -2, -3, and -4). The ectodomain of nectin family members comprises three Ig-like domains (V, C, C). Nectins localize at the adherens junctions (AJ) in epithelial and endothelial cells where they serve as adhesion molecules. Actin-based AJs play a role in mechanical adhesion, cellular morphogenesis and cellular differentiation. Nectin associates with the actin cytoskeleton through its interaction with the actin filament-binding protein afadin. Nectin 4 and afadin co-localize at cadherin-based adherens junctions in MDCKII epithelial cells. Nectin 4 and nectin 3 share a common binding region in the V domain of Nectin 1 and thus compete for Nectin 1 binding. The Nectin 3/4 binding domain maps to the C-C'-C''-D ? strands of the V domain of Nectin 1. Unlike other nectins, which are more widely expressed, Nectin 4 is mainly expressed in the placenta.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Angiogenesis, Cancer, Immuno Oncology