Recombinant Platelet endothelial cell adhesion molecule Antibody

Rabbit Monoclonal Antibody [Clone PECAM1/8594R]

Catalog No	Format	Size
5175-RBM46-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5175-RBM46-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5175-RBM46-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	PECAM1/8594R
Immunogen	Recombinant fragment (around aa625-738) of the human CD31 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	82.52kDa
Cellular Localization	Cell junction, Cell membrane, Membrane raft
Species Reactivity	Human
Positive Control	Jurkat or THP-1 cells. Human Spleen, Kidney, Heart, tonsil or angiosarcoma.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Platelet endothelial cell adhesion molecule Antibody



SDS-PAGE Analysis of Purified CD31 Recombinant Rabbit Monoclonal Antibody (PECAM1/8594R). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of Human Heart, Mouse Heart, Rat Heart, HamsterHeart, Guinea pig Heart and Rabbit Heart tissue lysates using CD31 Recombinant Rabbit Monoclonal Antibody (PECAM1/8594R).



Western Blot Analysis of THP-1, Human Kidney and Human Heart lysates using CD31 Recombinant Rabbit Monoclonal Antibody (PECAM1/8594R).

Specificity & Comments

CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 MAb reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.