

Recombinant SATB2 (Colorectal Carcinoma Marker) Antibody

Rabbit Monoclonal Antibody [Clone SATB2/12795R]

Catalog No	Format	Size
23314-RBM37-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
23314-RBM37-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
23314-RBM37-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	SATB2/12795R
Immunogen	Recombinant fragment (around aa710-733) of the human SATB2 (SATB homeobox 2) protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	82.55kDa
Cellular Localization	Nucleus matrix
Species Reactivity	Human
Positive Control	Colon: A strong nuclear staining should be seen in virtually all columnar epithelial cells. Brain, K-562

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant SATB2 (Colorectal Carcinoma Marker) Antibody



Western blot analysis of Brain tissue lysates of different species using SATB2 Recombinant Rabbit Monoclonal Antibody (SATB2/12795R).

Western blot analysis of Human brain and K-562 lysates using SATB2 Recombinant Rabbit Monoclonal Antibody (SATB2/12795R).

Specificity & Comments

SATB2 is a DNA binding protein that specifically binds nuclear matrix attachment regions. It is involved in transcription regulation and chromatin remodeling. SATB2 expression in colorectal carcinomas (CRC) is correlated with good prognosis and in laryngeal squamous cell carcinoma it functions as a tumor suppressor wherein loss of expression is positively correlated with high tumor grade and recurrence. Moreover, SATB2, in combination with CK20, could identify almost all CRC s. Upper gastrointestinal (GI) carcinomas and pancreatic ductal carcinomas are usually negative for SATB2, and ovarian carcinomas, lung adenocarcinomas, and adenocarcinomas from other origin are rarely positive for SATB2. Additionally, SATB2 antibody can identify neuroendocrine neoplasms of colon and rectum because SATB2 is usually negative in neuroendocrine neoplasms of the GI tract, pancreas, and lung. More recently, it has been reported that SATB2 is a sensitive marker for tumors with osteoblastic differentiation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
