

Recombinant PU.1 (SPI-1) (B-Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone PU1/13315R]

Catalog No	Format	Size
6688-RBM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6688-RBM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6688-RBM6-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	PU1/13315R
Immunogen	Recombinant full-length human PU.1 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	31.08kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	K-562, Raji, Ramos

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant PU.1 (SPI-1) (B-Cell Marker) Antibody



Western Blot Analysis of K-562, Raji and Ramos lysates using PU.1-Monospecific Recombinant Rabbit Monoclonal Antibody (PU1/13315R).

Specificity & Comments

PU.1 is a member of the ETS family of transcription factors and is important for normal B-cell development. It is expressed in the myeloid lineage and in immature as well as mature B-lymphocytes, with the exception of plasma cells. PU.1 is expressed in germinal center B-cells and mantle B-cells. Various lymphomas are also positive for this marker. It is essential during early B-cell differentiation. The absence of PU.1 results in total block of B-cell development at the pre-pro stage. PU.1 plays a key role in normal myeloid differentiation, and regulates the expression of immunoglobulin and other genes that are important for B-cell development.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.