

Recombinant BAP1 (BRCA1 Associated Protein 1) Antibody

Mouse Monoclonal Antibody [Clone rBAP1/13664]

Catalog No	Format	Size
8314-MSM56-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
8314-MSM56-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
8314-MSM56-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	rBAP1/13664
Immunogen	Recombinant fragment (around aa430-729) of the human BAP1 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	80.36kDa
Cellular Localization	Chromosome, Cytoplasm, Nucleus
Species Reactivity	Human
Positive Control	Jurkat or PC-3 or HepG2 cells. Human testis, ovarian or breast carcinoma.


*Optimal dilution for a specific application should be determined.

Product Images for Recombinant BAP1 (BRCA1 Associated Protein 1) Antibody




Formalin-fixed, paraffin-embedded human breast stained with BAP1 Recombinant Mouse Monoclonal Antibody (rBAP1/13664). Inset: PBS instead of primary antibody; secondary only negative control.

Purity: SDS-PAGE



SDS-PAGE Analysis of Purified BAP Confirmation of Purity and Integrity of Antibody.



Western blot analysis of HeLa and HEK293 cell lysates using BAP1 Recombinant Mouse Monoclonal Antibody (rBAP1/13664).

Specificity & Comments

The BAP1 gene belongs to the ubiquitin C-terminal hydrolase subfamily of de-ubiquitination enzymes that are involved in the removal of ubiquitin from proteins. The encoded enzyme binds to the breast cancer type 1 susceptibility protein (BRCA1) via the RING finger domain of the latter and acts as a tumor suppressor. In addition, the enzyme may be involved in regulation of transcription, regulation of cell cycle and growth, response to DNA damage and chromatin dynamics. Germ line mutations in this gene may be associated with tumor predisposition syndrome (TPDS), which involves increased risk of cancers including malignant mesothelioma, Uveal melanoma and cutaneous melanoma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.