

Recombinant MDM2 Antibody

Mouse Monoclonal Antibody [Clone rMDM2/13318]

Catalog No	Format	Size
4193-MSM28-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4193-MSM28-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4193-MSM28-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rMDM2/13318
Immunogen	Recombinant fragment (around aa154-167) of the human MDM2 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	55.23kDa
Cellular Localization	Cytoplasm, Nucleolus, Nucleoplasm, Nucleus
Species Reactivity	Human
Positive Control	Human Brain.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant MDM2 Antibody



SDS-PAGE Analysis of Purified MDM2 Recombinant Mouse Monoclonal Antibody (rMDM2/13318). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of Human Brain tissue lysate using MDM2 Recombinant Mouse Monoclonal Antibody (rMDM2/13318).

Specificity & Comments

MDM2 is a nuclear phosphoprotein that binds and inhibits transactivation by tumor protein p53. It can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. Overexpression of MDM2 can result in excessive inactivation of tumor protein p53, diminishing its tumor suppressor function. This protein also affects the cell cycle, apoptosis, and tumorigenesis through interactions with other proteins, including retinoblastoma 1 and ribosomal protein L5. Overexpression of MDM2 protein is detected in a variety of cancers.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
