

Recombinant CD44 / HCAM (Cancer Stem Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rHCAM/9967]

Catalog No	Format	Size
960-MSM30-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
960-MSM30-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
960-MSM30-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	rHCAM/9967
Immunogen	Recombinant full-length human CD44 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	81.54kDa
Cellular Localization	Cell membrane, Cell projection, Microvillus, Secreted
Species Reactivity	Human
Positive Control	Human Intestine. A549 or HeLa cells.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant CD44 / HCAM (Cancer Stem Cell Marker) Antibody



Western Blot Analysis of Human Intestine, A549 and HeLa lysates using CD44 Recombinant Mouse Monoclonal Antibody (rHCAM/9967).

Specificity & Comments

This antibody recognizes an epitope encoded by exon v3 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemopoiesis tissues. CD44 is a single gene with 20 exons, of which 10 are normally expressed to encode the basic CD44 (H-CAM) molecule. The additional 10 exons (v1 to v10) are only expressed by alternative splicing of the nuclear RNA. The expression of specific cell adhesion molecule CD44 splice variants has been reported to be associated with metastasis in certain human malignancies.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.