

## Recombinant Desmoglein-3 (Squamous Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone DSG3/13651R]

Catalog No	Format	Size
1830-RBM18-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1830-RBM18-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1830-RBM18-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


### Product Details

<b>Clone</b>	DSG3/13651R
<b>Immunogen</b>	Recombinant fragment (around aa379-491) of the human Desmoglein-3 protein (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	107.53kDa
<b>Cellular Localization</b>	Cell junction, Cell membrane, Cytoplasm, Desmosome, Tight junction
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	A431 cells. Lung squamous cell carcinoma.

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant Desmoglein-3 (Squamous Cell Marker) Antibody

Purity: SDS-PAGE



SDS-PAGE Analysis of Purified Desmoglein-3 Recombinant Rabbit Monoclonal Antibody (DSG3/13651R). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

Recognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell types. Currently, three desmoglein subfamily members are identified and all are members of the cadherin cell adhesion molecule superfamily.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.