

## Recombinant Folate Receptor Alpha (FRalpha) / FOLR1 Antibody

Rabbit Monoclonal Antibody [Clone FOLR1/13794R]

Catalog No	Format	Size
2348-RBM14-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2348-RBM14-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2348-RBM14-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


### Product Details

<b>Clone</b>	FOLR1/13794R
<b>Immunogen</b>	Synthetic human folate receptor alpha protein Cellular Localization: Cytoplasmic/cell surface
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	29.82kDa
<b>Cellular Localization</b>	Apical cell membrane, Basolateral cell membrane, Cell membrane, Clathrin-coated vesicle, Cytoplasmic vesicle, Endosome, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Kidney.

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant Folate Receptor Alpha (FRalpha) / FOLR1 Antibody

Purity: SDS-PAGE



SDS-PAGE under non-reducing (NR) and reducing (R) conditions. The gel was stained with Coomassie Blue. The purity of the protein is 96%.

SDS-PAGE Analysis of FOLR1 Recombinant Rabbit Monoclonal Antibody (FOLR1/13794R) of Purity and Integrity of Antibody.

### **Specificity & Comments**

Folate receptor alpha (FR?) is a membrane-bound protein that facilitates high-affinity folate transport, essential for cell metabolism, DNA synthesis, and repair, particularly in rapidly dividing cancer cells with increased folate demands. This receptor, encoded by the FOLR1 gene, employs a potocytosis mechanism that leads to both membranous and cytoplasmic localization and is selectively overexpressed in certain epithelial malignancies while remaining highly restricted in normal tissues. FR? expression correlates with tumor stage and grade, suggesting a role in tumor progression by enhancing folate uptake or generating regulatory signals. Its regulation is influenced by extracellular folate levels, homocysteine accumulation, hormones, transcription factors, and genetic mutations. Members of the folate receptor family share highly conserved sequences in the open reading frames but differ in amino acids in the 5' untranslated regions and as a consequence can differ in function and tissue expression. Given the selective overexpression of FR? in tumors and its association with aggressive cancer phenotypes, FR? is a valuable marker for studying tumor biology and progression, particularly in epithelial cancers.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---