

Recombinant PAX3 Antibody

Mouse Monoclonal Antibody [Clone PAX3/8425]


Catalog No	Format	Size
5077-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5077-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5077-MSM5-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details	
Clone	PAX3/8425
Immunogen	Recombinant fragment (around aa150-350) of human PAX3 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	52.97kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Human skin and skeletal muscle tissue.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant PAX3 Antibody



SDS-PAGE Analysis of PAX3 Mouse Monoclonal Antibody (PAX3/8425) of Purity and Integrity of Antibody.

Specificity & Comments

Pax genes contain paired domains that share strong homology to genes in *Drosophila* which are involved in programming early development. The product of the PAX3 gene is a DNA-binding protein expressed during early neurogenesis. Pax-3 is a protein containing both a paired domain and a paired-type homeodomain. During early neurogenesis, Pax-3 expression is limited to mitotic cells in the ventricular zone of the developing spinal cord and to distinct regions in the hindbrain, midbrain and diencephalon. In 10-12 day embryos, expression of Pax-3 is also seen in neural crest cells of the developing spinal ganglia, the craniofacial mesectoderm and in limb mesenchyme. Mutations in the MITF and Pax-3 genes, encoding transcription factors, are responsible for Waardenburg syndrome II (WSII) and WSI/WSIII, respectively.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246)
orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.