

Recombinant Tubulin beta 3 / TUBB3 (Neuronal Stem Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone TUBB3/13762R]

Catalog No	Format	Size
10381-RBM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10381-RBM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10381-RBM13-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details	
Clone	TUBB3/13762R
Immunogen	Recombinant full-length human TUBB3 (Tubulin beta 3 class III) protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	50.43kDa
Cellular Localization	Cell projection, Cytoplasm, Cytoskeleton, Filopodium, Growth cone, Lamellipodium
Species Reactivity	Human
Positive Control	Expression is primarily restricted to central and peripheral nervous system

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Tubulin beta 3 / TUBB3 (Neuronal Stem Cell Marker) Antibody

Specificity & Comments

Tubulin is the major constituent of microtubules, protein filaments consisting of alpha- and beta-tubulin heterodimers (PubMed:34996871, PubMed:38305685, PubMed:38609661). Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms (PubMed:34996871, PubMed:38305685, PubMed:38609661). Below the cap, alpha-beta tubulin heterodimers are in GDP-bound state, owing to GTPase activity of alpha-tubulin (PubMed:34996871, PubMed:38609661). TUBB3 plays a critical role in proper axon guidance and maintenance (PubMed:20074521). Binding of NTN1/Netrin-1 to its receptor UNC5C might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:28483977). Plays a role in dorsal root ganglion axon projection towards the spinal cord (PubMed:28483977).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.