# Recombinant Fibroblast growth factor receptor 2 Antibody

Rabbit Monoclonal Antibody [Clone FGFR2/13877R]

| Catalog No | Format | Size |
|---|---|---|
| 2263-RBM1-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 2263-RBM1-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 2263-RBM1-P1ABX | Purified Ab WITHOUT BSA or Azide at 1.0mg/ml | 100 ug |

| Applications | Tested Dillution | Note |
|---|---|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

## Product Details

| | |
|---|---|
| **Clone** | FGFR2/13877R |
| **Immunogen** | Recombinant fragment of the human FGFR2 protein (exact sequence is proprietary) |
| **Host** | Rabbit |
| **Clonality** | Monoclonal |
| **Isotype / Light Chain** | IgG / Kappa |
| **Mol. Weight of Antigen** | 92.02kDa |
| **Cellular Localization** | Cell membrane, Cytoplasmic vesicle, Golgi apparatus, Secreted |
| **Species Reactivity** | Human |

**Optimal dilution for a specific application should be determined.*

## Product Images for Recombinant Fibroblast growth factor receptor 2 Antibody

### Specificity & Comments

Tyrosine-protein kinase that acts as a cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation, migration and apoptosis, and in the regulation of embryonic development. Required for normal embryonic patterning, trophoblast function, limb bud development, lung morphogenesis, osteogenesis and skin development. Plays an essential role in the regulation of osteoblast differentiation, proliferation and apoptosis, and is required for normal skeleton development. Promotes cell proliferation in keratinocytes and immature osteoblasts, but promotes apoptosis in differentiated osteoblasts. Phosphorylates PLCG1, FRS2 and PAK4. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. FGFR2 signaling is down-regulated by ubiquitination, internalization and degradation. Mutations that lead to constitutive kinase activation or impair normal FGFR2 maturation, internalization and degradation lead to aberrant signaling. Over-expressed FGFR2 promotes activation of STAT1.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.