

Recombinant Interleukin-7 (IL-7) Antibody

Rabbit Monoclonal Antibody [Clone IL7/16922R]

Catalog No	Format	Size
3574-RBM24-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3574-RBM24-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3574-RBM24-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	IL7/16922R
Immunogen	Recombinant fragment (around aa45-164) of the human IL-7 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	20.19kDa
Cellular Localization	Secreted
Species Reactivity	Human

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Interleukin-7 (IL-7) Antibody

Specificity & Comments

Hematopoietic cytokine that plays an essential role in the development, expansion, and survival of naive and memory T-cells and B-cells thereby regulating the number of mature lymphocytes and maintaining lymphoid homeostasis (PubMed:25870237, PubMed:7527823). Mechanistically, exerts its biological effects through a receptor composed of IL7RA subunit and the cytokine receptor common subunit gamma/CSF2RG (PubMed:8128231). Binding to the receptor leads to activation of various kinases including JAK1 or JAK3 depending on the cell type and subsequently propagation of signals through activation of several downstream signaling pathways including the PI3K/Akt/mTOR or the JAK-STAT5 (PubMed:18523275, PubMed:20974963).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.