

Recombinant OVOL2 / CRE-BPa Antibody

Rabbit Monoclonal Antibody [Clone OVOL2/13818R]

Catalog No	Format	Size
58495-RBM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
58495-RBM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
58495-RBM11-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	OVOL2/13818R
Immunogen	Recombinant fragment (around aa133-275) of the human OVOL2 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	30.44kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Expressed in testis, ovary, heart and skeletal muscle (PubMed:12213202)

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant OVOL2 / CRE-BPa Antibody

Specificity & Comments

Zinc-finger transcription repressor factor (PubMed:19700410). Plays a critical role in maintaining the identity of epithelial lineages by suppressing epithelial-to mesenchymal transition (EMT) mainly through the repression of ZEB1, an EMT inducer (By similarity). Positively regulates neuronal differentiation (By similarity). Suppresses cell cycling and terminal differentiation of keratinocytes by directly repressing MYC and NOTCH1 (PubMed:19700410). Important for the correct development of primordial germ cells in embryos (By similarity). Plays dual functions in thermogenesis and adipogenesis to maintain energy balance. Essential for brown/beige adipose tissue-mediated thermogenesis, is necessary for the development of brown adipocytes. In white adipose tissues, limits adipogenesis by blocking CEBPA binding to its transcriptional targets and inhibiting its transcription factor activity (By similarity).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.