

Recombinant CCL23 / Myeloid Progenitor Inhibitory Factor 1 Antibody

Rabbit Monoclonal Antibody [Clone CCL23/13846R]

| Catalog No | Format | Size |
|-----------------|--|--------|
| 6368-RBM9-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 6368-RBM9-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 6368-RBM9-P1ABX | Purified Ab WITHOUT BSA or Azide at 1.0mg/ml | 100 ug |

| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details

| | |
|-------------------------------|---|
| Clone | CCL23/13846R |
| Immunogen | Recombinant fragment (around aa23-120) of the human CCL23 protein (exact sequence is proprietary) |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | 13.44kDa |
| Cellular Localization | Secreted |
| Species Reactivity | Human |
| Positive Control | High levels in adult lung, liver, skeletal muscle and pancreas |

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant CCL23 / Myeloid Progenitor Inhibitory Factor 1 Antibody

Specificity & Comments

Shows chemotactic activity for monocytes, resting T-lymphocytes, and neutrophils, but not for activated lymphocytes. Inhibits proliferation of myeloid progenitor cells in colony formation assays. This protein can bind heparin. Binds CCR1. CCL23(19-99), CCL23(22-99), CCL23(27-99), CCL23(30-99) are more potent chemoattractants than CCL23.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.