

## Recombinant Alpha-synuclein Antibody

Rabbit Monoclonal Antibody [Clone SNCA/9907R]

Catalog No	Format	Size
6622-RBM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6622-RBM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6622-RBM5-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
--------------	------------------	------

### Product Details

<b>Clone</b>	SNCA/9907R
<b>Immunogen</b>	Recombinant fragment (around aa53-87) of the human Synuclein, alpha protein (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	14.46kDa
<b>Cellular Localization</b>	Axon, Cell projection, Cytoplasm, Membrane, Nucleus, Secreted, Synapse
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Highly expressed in presynaptic terminals in the central nervous system

*\*Optimal dilution for a specific application should be determined.*

### Product Images for Recombinant Alpha-synuclein Antibody

#### Specificity & Comments

Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed:20798282, PubMed:26442590, PubMed:28288128, PubMed:30404828). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed:28288128, PubMed:30404828). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed:30404828). Also acts as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5 (PubMed:20798282). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed:20798282). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed:26442590).

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

#### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.