

Recombinant CD74 (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rCLIP/7265]

Catalog No	Format	Size
972-MSM15-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
972-MSM15-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
972-MSM15-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
--------------	------------------	------

Product Details

Clone	rCLIP/7265
Immunogen	Recombinant fragment (around aa1-296) (N-terminus) of human CD74 (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	33.52kDa
Cellular Localization	Cell membrane, Endoplasmic reticulum membrane, Endosome, Golgi apparatus, Late endosome, Lysosome, Secreted, trans-Golgi network
Species Reactivity	Human
Positive Control	Detected in urine (at protein level)

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant CD74 (B-Cell Marker) Antibody

Specificity & Comments

Plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to the endosomal/lysosomal system where the antigen processing and binding of antigenic peptides to MHC class II takes place. Serves as cell surface receptor for the cytokine MIF., Binds to the peptide-binding site of MHC class II alpha/beta heterodimers forming an alpha-beta-CLIP complex, thereby preventing the loading of antigenic peptides to the MHC class II complex until its release by HLA-DM in the endosome., Stabilizes the conformation of mature CTSL by binding to its active site and serving as a chaperone to help maintain a pool of mature enzyme in endocytic compartments and extracellular space of antigen-presenting cells (APCs). Has antiviral activity by stymieing the endosomal entry of Ebola virus and coronaviruses, including SARS-CoV-2 (PubMed:32855215). Disrupts cathepsin-mediated Ebola virus glycoprotein processing, which prevents viral fusion and entry. This antiviral activity is specific to p41 isoform (PubMed:32855215).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.