

Cadherin-5 Antibody

Mouse Monoclonal Antibody [Clone CDH5/14593]

Catalog No	Format	Size
1003-MSM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1003-MSM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1003-MSM6-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	CDH5/14593
Immunogen	Recombinant fragment (around aa 621-768) of human CDH5 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b
Mol. Weight of Antigen	87.53kDa
Cellular Localization	Adherens junction, Cell junction, Cell membrane, Cytoplasm
Species Reactivity	Human
Positive Control	Expressed in endothelial cells (at protein level) (PubMed:27338829)

*Optimal dilution for a specific application should be determined.

Product Images for Cadherin-5 Antibody

Specificity & Comments

Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed:21269602). This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed:10861224). Plays a role in coupling actin fibers to cell junctions in endothelial cells, via acting as a cell junctional complex anchor for AMOTL2 and MAG11 (By similarity). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B (PubMed:20332120). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction (PubMed:20332120). Associates with CTNND1/p120-catenin to control CADH5 endocytosis (By similarity).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.